



Retinal changes in preeclampsia

Pegah Rashidian ¹, Roxana Safari ², Seyyed Amirhossein Salehi ³ and Shaghayegh Karami ²

¹ Reproductive Health Research Center, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

² School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

³ Student Research Committee, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ABSTRACT

Background: The eye undergoes a series of physiological changes during pregnancy in response to metabolic and hormonal adaptation. Preeclampsia (PE) is the main cause of maternal morbidity and is associated with serious pregnancy-related complications. Characterized by hypertension and proteinuria, PE affects 2–8% of pregnancies globally. Previous studies have indicated that PE compromises retinal health. We examined the effects of PE on retinal structure and vasculature.

Methods: We conducted a literature review by searching PubMed/MEDLINE, Embase, and Scopus using terms pertaining to PE and the retina. The review included articles published between January 1, 1990, and May 30, 2024. The articles were selected based on their relevance to the topic. The key findings are summarized to offer a comprehensive overview of current knowledge, highlight the pathophysiology and manifestations of PE-related retinal changes, and propose clinical implications, diagnostic clues, and management strategies.

Results: PE is associated with visual disturbances caused by various retinal changes, including arteriolar narrowing, vasospasm, hemorrhages, exudates, serous retinal detachment, macular edema, retinal vein occlusion, and choroidal ischemia. These are mostly evident on fundus examination and frequently resolve postpartum. The underlying pathophysiology involves endothelial dysfunction, hypertension, inflammation, and coagulopathy. These retinal changes have immediate clinical implications and long-term risks, necessitating early detection, prompt multidisciplinary management, and close follow-up. Although most PE-associated retinal disturbances spontaneously resolve after the termination of pregnancy, one-third of patients are likely to experience long-lasting consequences.

Conclusions: Visual disturbances may precede and portend the onset of PE. Early detection and prompt management will mitigate fetal and maternal morbidity and mortality. Understanding the underlying pathophysiology of PE-related retinal manifestations is crucial for optimal management, because the majority of manifestations are reversible. Although retinal changes secondary to PE typically resolves postpartum, some studies demonstrated that women with PE may have a higher long-standing risk of ocular disorders. A multidisciplinary team approach involving obstetricians, ophthalmologists, and healthcare providers is essential for the immediate and long-term management of ocular consequences in pregnancy. Consequently, additional studies associated with robust methodologies are required to develop clinical guidelines.

KEYWORDS

preeclampsia, retina, pregnancies, hypertensive retinopathies, retinal detachments, machine learning, artificial intelligence, maternal health, retinal hemorrhages, postpartum

Correspondence: Pegah Rashidian, Reproductive Health Research Center, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.
Email: pegah.rashidian@rocketmail.com. ORCID iD: <https://orcid.org/0000-0002-9167-9172>

How to cite this article: Rashidian P, Safari R, Salehi SAH, Karami S. Retinal changes in preeclampsia. Med Hypothesis Discov Innov Optom. 2024 Summer; 5(2): 76-84. DOI: <https://doi.org/10.51329/mehdiptometry201>

Received: 04 July 2024; Accepted: 23 August 2024



Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.



INTRODUCTION

Multiple organs of the pregnant women are involved during pregnancy to provide an appropriate environment for embryo development and to prepare the mother for delivery [1]. For example, the cardiovascular, respiratory, gastrointestinal, hematologic, and visual systems are influenced throughout pregnancy [2-4]. Although these alterations during pregnancy are often normal, 5–20% of pregnancies are associated with pathological events [5].

The ocular system undergoes a series of physiological modifications during pregnancy in response to metabolic and hormonal adaptations [4, 6]. These include an increase in eyelid pigmentation, melasma, dry eye disease, changes in corneal thickness and curvature, myopic shift, and reduction in visual field [4, 7, 8]. However, a spectrum of pregnancy-induced etiologies can lead to pathological vision changes [9]. Numerous maternal vision disorders are attributable to the progression of pre-existing ocular impairments such as diabetic retinopathy and glaucoma [10, 11].

Nevertheless, other categories of ophthalmic impairments are believed to develop *de novo* during gestation. For example, systemic diseases such as disseminated intravascular coagulation (DIC), which may occur more frequently during pregnancy, can lead to ocular abnormalities. Moreover, pregnancy complications such as preeclampsia (PE) might cause ophthalmic pathologies leading to visual disturbances [8, 9, 12].

PE is one of the most common maternal health concerns, affecting 2–8% of pregnant women worldwide, and is responsible for approximately 500,000 fetal deaths and 46,000 maternal deaths globally per year [13-15]. PE is a hypertensive disorder of pregnancy, defined as the *de novo* onset of hypertension after the 20th week of gestation, accompanied by proteinuria, with or without end-organ damage, such as serous retinal detachment, acute kidney injury, thrombocytopenia, hemolysis, and liver or neurological disturbances [14-17].

The retina does not undergo any particular modification during normal pregnancy. However, this complex ocular structure is frequently impaired or damaged in pregnant women with PE [17, 18]. It is composed of neural tissue and supporting cells located in the posterior segment of the eye. The retina plays a notable role in processing the visual information received and conveying signals to the brain through the optic nerve [19, 20]. Owing to its high metabolic activity and oxygen demand, the retina possesses unique vasculature that supplies its requirements while maintaining transparency for light transmission [19]. The hypertensive component of PE can induce vasoconstriction in the retinal vasculature with subsequent vascular narrowing [21]. As retinal function relies on intact vasculature and adequate blood perfusion, disruptions caused by this hypertensive disorder can precipitate various pathologies that manifest as visual symptoms [17, 18]. We aimed to synthesize and classify the existing body of knowledge pertaining to the pathophysiology, diagnosis, and management of PE-induced retinal disorders. To achieve this objective, we conducted an in-depth review of the pertinent literature spanning three decades of research.

METHODS

For this narrative review, we explored the effects of PE on the retina through a literature search spanning 34 years using electronic databases, including PubMed/MEDLINE, EMBASE, and Scopus. The keywords used in the search strategy included "preeclampsia," "hypertensive disorders of pregnancy," "retinal health," "retinal vasculature," "retinopathy," "hemorrhage," "retinal detachment," "macular edema," "choroidal ischemia," and "treatment." The search was limited to articles published from January 1, 1990, to May 30, 2024, and included narrative reviews, systematic reviews, original research articles, case reports, and observational studies. The selected articles were assessed based on their relevance to the impact of PE on retinal structure and function, including retinal vascular changes, pathological findings, diagnostic methodologies, and therapeutic interventions. This narrative synthesis integrates the findings of diverse studies to provide a comprehensive overview of current information and the knowledge gaps in the field of PE-related retinal complications.

RESULTS and DISCUSSION

Retinal changes attributable to PE

PE can lead to various systemic complications, including those affecting the eyes. The retinal changes observed in PE are mainly attributable to alterations in the microcirculation caused by hypertension and endothelial dysfunction [18]. These changes can be identified through ophthalmological examination [4] and may include the following:

Arteriolar narrowing and vasospasm: Among the retinal changes that occur during PE, vasospasm can play a critical role in subsequent events. Vasospasm and arteriolar narrowing can be observed in the choroidal arterioles and central retinal or posterior ciliary arteries [22]. Arteriolar narrowing is the primary response of retinal vasculature to systemic arterial hypertension during PE [23]. Narrowing of the choroidal arterioles can ultimately lead to choroidal ischemia, resulting in increased vascular permeability. Consequently, fluid accumulation between the neurosensory retina and the retinal pigment epithelium causes retinal detachment [24]. Moreover, arteriolar narrowing is the most common finding on fundus examination among women with PE [25, 26].

Hemorrhage: Retinal hemorrhage is an important sign of systemic vascular disease. Etiologies of this condition comprise a vast array of diagnoses. However, diagnostic clues can help to differentiate between these conditions. PE is an urgent obstetric condition and an important cause of retinal hemorrhage. During PE, intraretinal hemorrhage, Elschnig's spots caused by choroidal ischemia, and serous retinal detachment can be observed [23, 27, 28]. These events typically occur bilaterally and vary in intensity. Furthermore, Roth's spots are a common sign of preeclamptic retinal hemorrhage [28].

Exudates: The presence of retinal hard exudates is an important sign of an underlying systemic disease involving the retina. Among these diseases, diabetes mellitus, hypertension, and PE should be considered [21, 29]. Retinal hard exudates are composed of lipid and proteinaceous material. These exudates occur because of leakage caused by an impaired blood–retinal barrier. Among other conditions, PE is one cause of this impairment [30].

Serous retinal detachment: Serous retinal detachment is a rare cause of vision loss during pregnancy, occurring in less than 1% of preeclamptic cases [31]. However, the incidence of this complication increases in the more harmful variant of PE known as “hemolysis, elevated liver enzyme levels, and low platelet levels” (HELLP) syndrome [32]. Choroidal ischemia occurs because of vasospasm, leading to subretinal leakage and retinal detachment [33]. This condition is usually bilateral, diagnosed after childbirth, and associated with primipara and/or cesarean delivery [34]. Surgical intervention is not appropriate for managing this condition because optimal treatment focuses on addressing the underlying cause. In the context of PE, management involves stabilization and expedited fetal delivery [17, 31]. Fortunately, this condition usually resolves and rarely causes permanent vision loss [17, 33].

Macular edema: Macular edema is caused by fluid accumulation in the macula and central retina, which are responsible for sharp central vision. The macular accumulation of fluid results from an imbalance between the mechanisms that regulate retinal fluid influx and efflux. This condition can have various etiologies, including diabetes mellitus, age-related macular degeneration, retinal vein occlusion, and PE [35]. During PE, funduscopy commonly reveals macular edema, which is often concurrent with serous retinal detachment and other retinal alterations [36]. Similar to retinal detachment, macular edema typically requires several weeks to resolve; however, periocular corticosteroid administration can accelerate resolution and expedite vision recovery [37].

Retinal vein occlusion: During pregnancy, the risk of thromboembolism increases, with greater likelihood in the third trimester and the postpartum period. This hypercoagulable state may induce retinal vein occlusion even in normal pregnancy [38]. By contrast, PE—and its fulminant form, HELLP syndrome—can cause retinal vein occlusion [39]. HELLP syndrome is a thrombotic microangiopathic vasculopathy that can lead to coagulation in different parts of the vascular system, and in this context, can cause central retinal vein occlusion. Retinal vein occlusion secondary to PE is expected to occur during the postpartum period [39].

Choroidal ischemia and infarction: Mild arteriolar spasm in the bulbar conjunctival vessels is considered normal in typical pregnancies [40]. However, in PE, choroidal ischemia—and in severe cases, choroidal infarction—typically arises from pronounced arteriolar vasospasm. Serous retinal detachment may also result from choroidal ischemia [34]. In addition to retinal detachment, yellowish, opaque retinal pigment epithelial lesions, also known as Elschnig’s spots, have been mentioned as a manifestation of choroidal ischemia [41].

Pathophysiology

Endothelial dysfunction: PE initially leads to systemic endothelial dysfunction, which is central to its pathophysiology. During the early stages of pregnancy, PE begins with the aberrant invasion of trophoblasts during placenta formation, resulting in oxidative stress and reduced blood flow to the uterus and placenta [42]. This process increases production of anti-angiogenic factors, such as soluble fms-like tyrosine kinase 1 (sFlt-1) and soluble endoglin, and reduced production of placental growth factor, inducing an imbalance between pro- and anti-angiogenic factors [42, 43]. Increased concentrations of sFlt-1 counteract the effects of vascular endothelial growth factor (VEGF), resulting in endothelial cell damage and impaired vascular healing mechanisms [42]. Increased VEGF levels stimulate the release of nitric oxide (NO) by increasing the expression of NO synthase in endothelial cells [21, 44]. This activates endothelial cells, causing changes in capillary structure and a decrease in NO levels [45]. NO stimulates the activation of soluble guanylate cyclase, which in turn triggers the creation of cyclic guanosine monophosphate (cGMP) [46]. Additionally, in PE, the balance between vasoconstrictive and vasodilatory prostaglandins is disrupted, leading to reduced levels of prostaglandin I₂ [47, 48]. Prostaglandin I₂ stimulates adenylyl cyclase and enhances the production of cyclic adenosine monophosphate (cAMP). Both cGMP and cAMP reduce the levels of calcium within cells, resulting in the relaxation of smooth muscle and subsequent vasodilation [49–51]. Therefore, endothelial dysfunction results in notable alterations in retinal structure and function, including thickening of the capillary basement membrane, loss of perivascular cells, impairment of the blood–retinal barrier, and neovascularization [52].

Hypertension: Hypertensive retinopathy refers to the retinal vascular alterations caused by high systemic blood pressure. In the early stages, vasospasm and elevated vascular tone cause generalized narrowing of the retinal arterioles and arteries. This response to elevated blood pressure results in localized or widespread constriction of blood vessels. Furthermore, vascular leakage is caused by elevated blood vessel permeability [53]. The resultant retinal changes include a reduction in the retinal artery-to-vein ratio and the presence of cotton wool spots, Elschnig’s spots, and serous retinal detachments [43, 54, 55]. In chronic hypertension, structural changes are evident in the vessel wall, including thickening of the inner layer and degradation of hyaline [56]. Microaneurysms may also develop under these circumstances. Cotton wool spots are ischemic foci in the retinal nerve fiber layer caused by severe hypertension [56]. Severe hypertension can lead to elevated intracranial pressure, compromised optic nerve blood supply, and severe bilateral optic disc swelling [45].

Coagulopathy: The hypercoagulable state caused by PE also exacerbates retinal damage. Elevated concentrations of procoagulant substances and reduced thrombolysis lead to the development of microthrombi within the retinal blood vessels. Microthrombi can induce retinal ischemia and infarction [55], resulting in retinal damage manifesting as cotton wool spots and retinal hemorrhages [36]. Pregnancy-related hypercoagulable states may arise because of the increased production of thrombin and procoagulant factors, with reduced levels of endogenous anticoagulants such as protein C, protein S, and antithrombin III [57].

Furthermore, individuals with PE exhibit increased levels of tissue factor, factor VIII, von Willebrand factor antigen, and fibrinopeptides A and B [21, 51, 58].

Inflammation: PE contributes to retinal injury through the activation of various inflammatory factors. Elevated levels of cytokines, such as interleukin-6 and tumor necrosis factor-alpha, lead to increased vascular permeability and leukocyte adhesion to the endothelium, exacerbating endothelial damage and retinal manifestations, including hemorrhages and exudates [59, 60]. Oxidative stress is crucial in the development of PE and affects the retina. The imbalance between pro- and antioxidant systems leads to the build-up of reactive oxygen species, which directly damage retinal cells and exacerbate endothelial dysfunction. Additionally, oxidative stress stimulates inflammation, thereby exacerbating vascular injury and causing retinal damage [43, 61]. Figure 1 summarizes the retinal changes associated with PE and their underlying pathophysiological mechanisms, providing a clear overview of the relationships between these conditions.




Figure 1. Retinal changes and pathophysiological mechanisms in preeclampsia

Clinical implications and diagnosis

PE can cause retinal damage, resulting in a spectrum of visual complaints that correlate with the severity of retinal injury. Visual symptoms manifest in approximately 40% of women with severe PE [36]. Patients with PE commonly have a range of ocular symptoms, including photopsia, blurred vision, color vision changes, blindness, visual field abnormalities, serous retinal detachment, and cortical blindness [21, 66]. Blurred vision is a prevalent symptom that can arise from retinal edema, ischemia, or hemorrhages [21, 67]. Cortical blindness is a severe and alarming consequence of PE that may resolve following the end of pregnancy, often within 8 days. It can also occur because of vasospasm and increased permeability of blood vessels of the brain [21, 58]. Neuroimaging may reveal a range of findings in the occipital lobes [58].

Photopsia is a sudden visual disturbance manifesting as flashes of light in the field of vision, commonly attributable to retinal detachment [68]. Serous retinal detachment occurs when the neurosensory retina separates from the retinal pigment epithelium, leading to vision impairment in patients with PE. Although it can manifest at any time during pregnancy, it typically occurs just before or shortly after childbirth. Bullous and bilateral detachment are common manifestations [69], with most patients experiencing sudden vision loss. Patients with the most severe types of PE exhibit the most unfavorable fundoscopic manifestations. Some studies have indicated an unfavorable fetal prognosis, whereas others have found no prognostic consequences for the fetus [21, 23].

The diagnosis of hypertensive retinopathy requires a thorough ophthalmic examination, including fundus photography [70], B-scan ultrasonography [71], and optical coherence tomography (OCT) [72]. These diagnostic tools can be used effectively in pregnant women. These noninvasive modalities are crucial for assessing retinal injuries, optic nerve changes, and choroidal abnormalities. Typically, invasive procedures, such as fundus fluorescein angiography, are avoided because of uncertainty regarding their teratogenic effects [36, 73]. Fundus photography captures high-resolution images of the retina, allowing documentation of hypertensive retinal alterations [70, 74]. OCT is another valuable tool for evaluating retinal thickness and identifying macular edema, which can develop in severe cases of hypertensive retinopathy [72]. OCT angiography facilitates detailed blood vessel image acquisition in different retinal layers, providing a three-dimensional representation of the vascular layers [62, 67]. Furthermore, a grading system designated as the Keith–Wagener–Barker classification is employed to evaluate the severity of hypertensive retinopathy. This approach classifies hypertensive retinopathy into four grades based on the severity of retinal arteriolar alterations, hemorrhages, exudates, and optic disc swelling [54, 75].

Management

Although most PE-associated retinal disturbances spontaneously resolve after the termination of pregnancy, one-third of patients are likely to experience long-lasting consequences [17, 18].

The first step in managing the ocular complications of PE is elimination of the underlying cause [21]. Delivery is the primary treatment modality for PE. The termination of gestation reverses most ophthalmic disturbances [17, 32, 76]. However, several existing complementary remedies restrict the progression of PE to eclampsia and prevent further sequelae [76].

Elevated blood pressure associated with PE requires the administration of antihypertensive medications [77]. The American College of Obstetricians and Gynecologists (ACOG) recommends lowering the blood pressure to less than 160/110 mmHg [78]. Because of contraindications and other concerns of medication use during pregnancy, the choice of antihypertensive agents is limited. Labetalol, hydralazine, methyldopa, beta-blockers, nifedipine, diazoxide, and nitroprusside are currently administered to control hypertension [76, 77]. Methyldopa is the currently preferred medication and is orally administered in divided doses [79]. Labetalol can be administered orally or intravenously [77]. Other medications are used less frequently as monotherapy or in combination regimens [76, 78].

The second most notable agent in PE treatment is magnesium sulfate [17, 79]. This neuroprotective substance prevents the progression of PE to eclampsia [17, 79]. The ACOG exclusively recommends magnesium sulfate administration in severe cases of PE [79]. Studies have also illustrated the effects of nutritional factors and exercise on the control of PE. Aerobic exercise combined with a low-salt, antioxidant-rich diet is beneficial for pregnant women and can prevent and control hypertensive disorders [77].

In addition to general PE management and conservative approaches, some retinal injuries may require specific treatments and interventions. Depending on the retinal pathology, supplemental ophthalmic medications and interventions can be cautiously employed to prioritize the well-being of the mother and fetus [4, 55, 80, 81].

The use of medications during pregnancy is subject to numerous restrictions, as several ophthalmic agents confer potential risks to developing embryos. Notably, acetazolamide and certain subtypes of beta-blockers have been identified as detrimental to the fetus [55, 82, 83]. However, the limited available evidence suggests no teratogenic effects of topical prostaglandin analogs [55, 84]. In the absence of contraindications, preferred ophthalmic agents can be prescribed according to the circumstances and ophthalmologist's diagnosis [55].

Although most cases of retinal injury and detachment recover after delivery, surgery may rarely be required. Surgeries and invasive procedures must be considered only in urgent circumstances, and preferably in the presence of the obstetrical team [55, 81].

Finally, pregnant women experiencing ocular manifestations of PE benefit from the collaboration of a multidisciplinary team comprising primary eye health care providers, ophthalmologists, and gynecologists aimed at optimizing maternal outcomes while highlighting the significance of postpartum follow-up care [4, 8, 85].

This narrative review summarizes key issues concerning PE-related retinal manifestations, pathophysiology, diagnosis, and management, according to studies published over a time span of more than three decades, as a road map for health care providers. However, the lack of a comprehensive systematic search strategy and meta-analysis is a limitation of this study. Further studies focusing on certain ocular manifestations of PE and their long-term consequences, preferably using a meta-analytic approach, could clarify the precise likelihood of visual consequences. In addition, investigating the association between observed ocular changes and maternal/fetal morbidity and mortality could provide additional practical guidance on this topic. Early prediction of PE is vital for controlling disease progression. Recent studies have exploited machine learning and artificial intelligence (AI) in the risk prediction for PE and related subtypes [86]. The use of AI algorithm-based retinal fundus photographs has demonstrated value in predicting PE [87]. Further studies using similar approaches could develop AI-based applications to predict short- and long-term visual outcomes in women who develop PE and its related complications.

CONCLUSIONS

PE, recognized as one of the most prevalent and severe complications of pregnancy, may compromise retinal integrity by disrupting retinal vascular structure and blood perfusion. Investigation of the impact of PE on retinal health has highlighted the critical need for early detection and effective and timely management of this condition throughout gestation. Visual disturbances may precede and portend the onset of PE. Therefore, early detection and prompt management will mitigate fetal and maternal morbidity and mortality. These findings emphasize the associations between PE and diverse retinal manifestations, thereby accentuating the potential long-term implications for maternal visual health. Future efforts should prioritize sustained interdisciplinary collaboration between obstetricians and ophthalmologists to advance our understanding of these connections and to devise targeted interventions aimed at maternal ocular well-being.

ETHICAL DECLARATIONS

Ethical approval: Not required.

Conflict of interest: None.

FUNDING

None.

ACKNOWLEDGMENTS

None.

REFERENCES

1. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr.* 2016 Mar-Apr;27(2):89-94. doi: 10.5830/CVJA-2016-021. PMID: 27213856; PMCID: PMC4928162.
2. Kazma JM, van den Anker J, Allegaert K, Dallmann A, Ahmadzia HK. Anatomical and physiological alterations of pregnancy. *J Pharmacokinet Pharmacodyn.* 2020 Aug;47(4):271-285. doi: 10.1007/s10928-020-09677-1. Epub 2020 Feb 6. PMID: 32026239; PMCID: PMC7416543.
3. Bossung V, Singer A, Ratz T, Rothenbühler M, Leeners B, Kimmich N. Changes in Heart Rate, Heart Rate Variability, Breathing Rate, and Skin Temperature throughout Pregnancy and the Impact of Emotions-A Longitudinal Evaluation Using a Sensor Bracelet. *Sensors (Basel).* 2023 Jul 23;23(14):6620. doi: 10.3390/s23146620. PMID: 37514915; PMCID: PMC10385491.
4. Morya AK, Gogia S, Gupta A, Prakash S, Solanki K, Naidu AD. Motherhood: What every ophthalmologist needs to know. *Indian J Ophthalmol.* 2020 Aug;68(8):1526-1532. doi: 10.4103/ijo.IJO_2033_19. PMID: 32709768; PMCID: PMC7640830.
5. Çelik FP, Güneri SE. The Relationship between Adaptation to Pregnancy and Prenatal Attachment in High-Risk Pregnancies. *Psychiatr Danub.* 2020 Nov;32(Suppl 4):568-575. PMID: 33212465.
6. Naderan M. Ocular changes during pregnancy. *J Curr Ophthalmol.* 2018 Jan 3;30(3):202-210. doi: 10.1016/j.joco.2017.11.012. PMID: 30197948; PMCID: PMC6127369.
7. Yenerel NM, Küçümen RB. Pregnancy and the Eye. *Turk J Ophthalmol.* 2015 Oct;45(5):213-219. doi: 10.4274/tjo.43815. Epub 2015 Oct 5. PMID: 27800235; PMCID: PMC5082244.
8. Qin Q, Chen C, Cugati S. Ophthalmic associations in pregnancy. *Aust J Gen Pract.* 2020 Oct;49(10):673-680. doi: 10.31128/AJGP-10-19-5113. PMID: 33015686.
9. Anton N, Doroftei B, Ilie OD, Ciuntu RE, Bogdănici CM, Nechita-Dumitriu I. A Narrative Review of the Complex Relationship between Pregnancy and Eye Changes. *Diagnostics (Basel).* 2021 Jul 23;11(8):1329. doi: 10.3390/diagnostics11081329. PMID: 34441264; PMCID: PMC8394444.
10. Rashidian P. Pregnancy and diabetic retinopathy. *Med Hypothesis Discov Innov Optom.* 2024 Spring; 5(1): 35-42. doi:10.51329/mehdiophtometry195
11. Salim S. Glaucoma in pregnancy. *Curr Opin Ophthalmol.* 2014 Mar;25(2):93-7. doi: 10.1097/ICU.000000000000029. PMID: 24469077.
12. Rzeszutarska A, Szczapa-Jagustyn J, Kociecki J. Ophthalmological problems in pregnancy - a review. *Ginekol Pol.* 2020;91(8):473-477. doi: 10.5603/GP.2020.0080. PMID: 32902845.

13. Yang Y, Le Ray I, Zhu J, Zhang J, Hua J, Reilly M. Preeclampsia Prevalence, Risk Factors, and Pregnancy Outcomes in Sweden and China. *JAMA Netw Open*. 2021 May 3;4(5):e218401. doi: 10.1001/jamanetworkopen.2021.8401. PMID: 33970258; PMCID: PMC8111481.
14. Ives CW, Sinkey R, Rajapreyar I, Tita ATN, Oparil S. Preeclampsia-Pathophysiology and Clinical Presentations: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2020 Oct 6;76(14):1690-1702. doi: 10.1016/j.jacc.2020.08.014. PMID: 33004135.
15. Hackelöer M, Schmidt L, Verlohren S. New advances in prediction and surveillance of preeclampsia: role of machine learning approaches and remote monitoring. *Arch Gynecol Obstet*. 2023 Dec;308(6):1663-1677. doi: 10.1007/s00404-022-06864-y. Epub 2022 Dec 25. PMID: 36566477; PMCID: PMC9790089.
16. Mészáros B, Kukor Z, Valent S. Recent Advances in the Prevention and Screening of Preeclampsia. *J Clin Med*. 2023 Sep 17;12(18):6020. doi: 10.3390/jcm12186020. PMID: 37762960; PMCID: PMC10532380.
17. Soullane S, Rhéaume MA, Auger N. Preeclampsia and the Retina. *Curr Hypertens Rep*. 2024 Apr;26(4):169-174. doi: 10.1007/s11906-023-01290-0. Epub 2023 Dec 22. PMID: 38133842.
18. He X, Ji Y, Yu M, Tong Y. Choriorretinal Alterations Induced by Preeclampsia. *J Ophthalmol*. 2021 Mar 10;2021:8847001. doi: 10.1155/2021/8847001. PMID: 33777446; PMCID: PMC7969093.
19. Wright WS, Eshaq RS, Lee M, Kaur G, Harris NR. Retinal Physiology and Circulation: Effect of Diabetes. *Compr Physiol*. 2020 Jul 8;10(3):933-974. doi: 10.1002/cphy.c190021. PMID: 32941691; PMCID: PMC10088460.
20. Pfitz M, Bleau M, Bouskila J. The Retina: A Window into the Brain. *Cells*. 2021 Nov 23;10(12):3269. doi: 10.3390/cells10123269. PMID: 34943777; PMCID: PMC8699497.
21. Abu Samra K. The eye and visual system in the preeclampsia/eclampsia syndrome: What to expect? *Saudi J Ophthalmol*. 2013 Jan;27(1):51-3. doi: 10.1016/j.sjopt.2012.04.003. Epub 2012 Apr 23. PMID: 23964188; PMCID: PMC3729391.
22. Maslovitz S, Lessing JB, Kupferminc MJ. Bilateral retinal detachment in preeclamptic women with thrombophilia. *Int J Gynaecol Obstet*. 2005 Oct;91(1):65-6. doi: 10.1016/j.ijgo.2005.06.016. PMID: 16098529.
23. Gupta A, Kaliaperumal S, Setia S, Suchi ST, Rao VA. Retinopathy in preeclampsia: association with birth weight and uric acid level. *Retina*. 2008 Oct;28(8):1104-10. doi: 10.1097/IAE.0b013e3181744122. PMID: 18779717.
24. Roos NM, Wiegman MJ, Jansonius NM, Zeeman GG. Visual disturbances in (pre)eclampsia. *Obstet Gynecol Surv*. 2012 Apr;67(4):242-50. doi: 10.1097/OGX.0b013e318250a457. PMID: 22495060.
25. Warad C, Midha B, Pandey U, Sivakrishna P, Jain A, Bagadia B, Makhija V, Pravin Patil B, Cheguri S, B K B. Ocular Manifestations in Pregnancy-Induced Hypertension at a Tertiary Level Hospital in Karnataka, India. *Cureus*. 2023 Feb 12;15(2):e34887. doi: 10.7759/cureus.34887. PMID: 36925976; PMCID: PMC10011940.
26. Chandran JR, Narayanan IB, Rajan J. Ocular Manifestations: Are They Significant in Hypertensive Disorders of Pregnancy? *J Obstet Gynaecol India*. 2021 Apr;71(2):118-123. doi: 10.1007/s13224-020-01385-7. Epub 2020 Nov 17. PMID: 34149212; PMCID: PMC8167031.
27. Kaur B, Taylor D. Retinal haemorrhages. *Arch Dis Child*. 1990 Dec;65(12):1369-72. doi: 10.1136/adc.65.12.1369. PMID: 2103739; PMCID: PMC1793090.
28. Kanukollu VM, Ahmad SS. Retinal Hemorrhage. 2023 Aug 8. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: 32809612.
29. Saito Y. [Retinchoroidal changes in toxemia of pregnancy with the relation to hypertensive retinopathy and choroidopathy]. *Nippon Ganka Gakkai Zasshi*. 1990 Aug;94(8):748-55. Japanese. PMID: 2239551.
30. Davoudi S, Papavasileiou E, Roohipoor R, Cho H, Kudrimoti S, Hancock H, Hoadley S, Andreoli C, Husain D, James M, Penman A, Chen CJ, Sobrin L. OPTICAL COHERENCE TOMOGRAPHY CHARACTERISTICS OF MACULAR EDEMA AND HARD EXUDATES AND THEIR ASSOCIATION WITH LIPID SERUM LEVELS IN TYPE 2 DIABETES. *Retina*. 2016 Sep;36(9):1622-9. doi: 10.1097/IAE.0000000000001022. PMID: 26991647; PMCID: PMC4993650.
31. Hussain SA, O'Shea BJ, Thagard AS. Preeclamptic serous retinal detachment without hypertension: A case report. *Case Rep Womens Health*. 2019 Jan 23;21:e00098. doi: 10.1016/j.crwh.2019.e00098. PMID: 30733925; PMCID: PMC6358546.
32. Radha Bai Prabhu T. Serious Visual (Ocular) Complications in Pre-eclampsia and Eclampsia. *J Obstet Gynaecol India*. 2017 Oct;67(5):343-348. doi: 10.1007/s13224-017-0975-6. Epub 2017 Mar 10. PMID: 28867885; PMCID: PMC5561762.
33. Phang DSK, Ariffin N, Abd Aziz H, Vendargon FM, Sonny Teo KS. Bilateral Serous Retinal Detachment in Pregnancy. *Cureus*. 2022 Oct 7;14(10):e30019. doi: 10.7759/cureus.30019. PMID: 36348857; PMCID: PMC9637277.
34. Raposo JTBV, Melo BCDS, Maciel NFBB, Leite SD, Rebelo ÓRC, Lima AMF. Serous Retinal Detachment in Pre-eclampsia: Case Report and Literature Review. *Rev Bras Ginecol Obstet*. 2020 Nov;42(11):772-773. doi: 10.1055/s-0040-1718448. Epub 2020 Nov 30. PMID: 33254274; PMCID: PMC10309245.
35. Kohli P, Tripathy K, Patel BC. Macular Edema. 2024 Apr 24. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: 35015421.
36. Stern EM, Blace N. Ophthalmic Pathology of Preeclampsia. 2022 Oct 24. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: 35015414.
37. Mansour AM, Medawar WA. Periocular corticosteroid in malignant hypertensive and preeclamptic choroidopathy. *Retin Cases Brief Rep*. 2009 Spring;3(2):144-6. doi: 10.1097/ICB.0b013e318162d998. PMID: 25391058.
38. Eldor A. Thrombophilia and its treatment in pregnancy. *J Thromb Thrombolysis*. 2001 Sep;12(1):23-30. doi: 10.1023/a:1012730325902. PMID: 11711685.
39. Gonzalvo FJ, Abecia E, Pinilla I, Izaguirre LB, Oliván JM, Honrubia FM. Central retinal vein occlusion and HELLP syndrome. *Acta Ophthalmol Scand*. 2000 Oct;78(5):596-8. doi: 10.1034/j.1600-0420.2000.078005596.x. PMID: 11037923.
40. Lee C, Hsu TY, Ou CY, Chang SY, Soong YK. Retinal detachment in postpartum preeclampsia and eclampsia: report of two cases. *Changcheng Yi Xue Za Zhi*. 1999 Sep;22(3):520-4. PMID: 10584429.
41. Errera MH, Kohly RP, da Cruz L. Pregnancy-associated retinal diseases and their management. *Surv Ophthalmol*. 2013 Mar-Apr;58(2):127-42. doi: 10.1016/j.survophthal.2012.08.001. PMID: 23410822.
42. Tomimatsu T, Mimura K, Matsuzaki S, Endo M, Kumasawa K, Kimura T. Preeclampsia: Maternal Systemic Vascular Disorder Caused by Generalized Endothelial Dysfunction Due to Placental Antiangiogenic Factors. *Int J Mol Sci*. 2019 Aug 30;20(17):4246. doi: 10.3390/ijms20174246. PMID: 31480243; PMCID: PMC6747625.

43. Aouache R, Biquard L, Vaiman D, Miralles F. Oxidative Stress in Preeclampsia and Placental Diseases. *Int J Mol Sci*. 2018 May 17;19(5):1496. doi: [10.3390/ijms19051496](#). PMID: [29772777](#); PMCID: [PMC5983711](#).
44. Bouloumié A, Schini-Kerth VB, Busse R. Vascular endothelial growth factor up-regulates nitric oxide synthase expression in endothelial cells. *Cardiovasc Res*. 1999 Mar;41(3):773-80. doi: [10.1016/s0008-6363\(98\)00228-4](#). PMID: [10435050](#).
45. Henderson AD, Bruce BB, Newman NJ, Biousse V. Hypertension-related eye abnormalities and the risk of stroke. *Rev Neurol Dis*. 2011;8(1-2):1-9. PMID: [21769065](#); PMCID: [PMC3448945](#).
46. Wittenborn EC, Marletta MA. Structural Perspectives on the Mechanism of Soluble Guanylate Cyclase Activation. *Int J Mol Sci*. 2021 May 21;22(11):5439. doi: [10.3390/ijms22115439](#). PMID: [34064029](#); PMCID: [PMC8196705](#).
47. Atallah A, Lecarpentier E, Goffinet F, Doret-Dion M, Gaucherand P, Tsatsaris V. Aspirin for Prevention of Preeclampsia. *Drugs*. 2017 Nov;77(17):1819-1831. doi: [10.1007/s40265-017-0823-0](#). PMID: [29039130](#); PMCID: [PMC5681618](#).
48. Walsh SW. Eicosanoids in preeclampsia. *Prostaglandins Leukot Essent Fatty Acids*. 2004 Feb;70(2):223-32. doi: [10.1016/j.plefa.2003.04.010](#). PMID: [14683695](#).
49. Robinson ES, Khankin EV, Karumanchi SA, Humphreys BD. Hypertension induced by vascular endothelial growth factor signaling pathway inhibition: mechanisms and potential use as a biomarker. *Semin Nephrol*. 2010 Nov;30(6):591-601. doi: [10.1016/j.semnephrol.2010.09.007](#). PMID: [21146124](#); PMCID: [PMC3058726](#).
50. Makhoul S, Walter E, Pagel O, Walter U, Sickmann A, Gambaryan S, Smolenski A, Zahedi RP, Jurk K. Effects of the NO/soluble guanylate cyclase/cGMP system on the functions of human platelets. *Nitric Oxide*. 2018 Jun 1;76:71-80. doi: [10.1016/j.niox.2018.03.008](#). Epub 2018 Mar 14. PMID: [29550521](#).
51. Kontovazainitis CG, Gialampirinou D, Theodoridis T, Mitsiakos G. Hemostasis in Pre-Eclamptic Women and Their Offspring: Current Knowledge and Hemostasis Assessment with Viscoelastic Tests. *Diagnostics (Basel)*. 2024 Feb 5;14(3):347. doi: [10.3390/diagnostics14030347](#). PMID: [38337863](#); PMCID: [PMC10855316](#).
52. Sorrentino FS, Matteini S, Bonifazzi C, Sebastiani A, Parmeggiani F. Diabetic retinopathy and endothelin system: microangiopathy versus endothelial dysfunction. *Eye (Lond)*. 2018 Jul;32(7):1157-1163. doi: [10.1038/s41433-018-0032-4](#). Epub 2018 Mar 9. PMID: [29520046](#); PMCID: [PMC6043602](#).
53. Modi P, Arsiwalla T. Hypertensive Retinopathy. 2023 Jul 4. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: [30252236](#).
54. Tsukikawa M, Stacey AW. A Review of Hypertensive Retinopathy and Choriorretinopathy. *Clin Optom (Auckl)*. 2020 May 5;12:67-73. doi: [10.2147/OPTO.S183492](#). PMID: [32440245](#); PMCID: [PMC7211319](#).
55. Rosenthal JM, Johnson MW. Management of Retinal Diseases in Pregnant Patients. *J Ophthalmic Vis Res*. 2018 Jan-Mar;13(1):62-65. doi: [10.4103/jovr.jovr_195_17](#). PMID: [29403592](#); PMCID: [PMC5782459](#).
56. Schmidt D. The mystery of cotton-wool spots - a review of recent and historical descriptions. *Eur J Med Res*. 2008 Jun 24;13(6):231-66. PMID: [18558551](#).
57. Bahar MM, Ghalandarpoor-Attar SN, Shabani A, Hantoushadeh S, Tabatabaei SA, Ghalandarpoor-Attar SM. Idiopathic combined retinal vessels occlusion in a pregnant woman: a case report. *J Med Case Rep*. 2022 May 15;16(1):191. doi: [10.1186/s13256-022-03421-8](#). PMID: [35568885](#); PMCID: [PMC9107689](#).
58. Nagy ZZ. Review of the ophthalmic symptoms of preeclampsia. *Developments in Health Sciences*. 2020 Oct 31;3(1):21-3. doi: [10.1556/2066.2020.00005](#).
59. Wang Y, Gu Y, Alexander JS, Lewis DF. Preeclampsia Status Controls Interleukin-6 and Soluble IL-6 Receptor Release from Neutrophils and Endothelial Cells: Relevance to Increased Inflammatory Responses. *Pathophysiology*. 2021 Apr 8;28(2):202-211. doi: [10.3390/pathophysiology28020013](#). PMID: [35366257](#); PMCID: [PMC8830466](#).
60. Swellam M, Samy N, Wahab SA, Ibrahim MS. Emerging role of endothelial and inflammatory markers in preeclampsia. *Dis Markers*. 2009;26(3):127-33. doi: [10.3233/DMA-2009-0622](#). PMID: [19597295](#); PMCID: [PMC3833325](#).
61. Chiarello DI, Abad C, Rojas D, Toledo F, Vázquez CM, Mate A, Sobrevia L, Marín R. Oxidative stress: Normal pregnancy versus preeclampsia. *Biochim Biophys Acta Mol Basis Dis*. 2020 Feb 1;1866(2):165354. doi: [10.1016/j.bbdis.2018.12.005](#). Epub 2018 Dec 24. PMID: [30590104](#).
62. Di Marco E, Aiello F, Lombardo M, Di Marino M, Missiroli F, Mancino R, Ricci F, Nucci C, Noce A, Di Daniele N, Cesareo M. A literature review of hypertensive retinopathy: systemic correlations and new technologies. *Eur Rev Med Pharmacol Sci*. 2022 Sep;26(18):6424-6443. doi: [10.26355/eurrev_202209_29742](#). PMID: [36196693](#).
63. Iqra HH, Anggraini MA, Junior DG, Andari MY, Danianto A. Serous retinal detachment in pre-eclampsia: a systematic review of case report. *Jurnal Kedokteran Brawijaya*. 2023 May 8;198-204. doi: [10.21776/ub.jkb.2023.032.03.11](#).
64. Kim IK, Shin JE, Kim MJ, Ra H, Baek J. Quantitative analysis of choroidal morphology in preeclampsia during pregnancy according to retinal change. *Sci Rep*. 2023 Aug 14;13(1):13171. doi: [10.1038/s41598-023-40144-2](#). PMID: [37580383](#); PMCID: [PMC10425443](#).
65. Kirolos S, Skilton M, Patel S, Amott C. A Systematic Review of Vascular Structure and Function in Pre-eclampsia: Non-invasive Assessment and Mechanistic Links. *Front Cardiovasc Med*. 2019 Nov 15;6:166. doi: [10.3389/fcvm.2019.00166](#). PMID: [31803759](#); PMCID: [PMC6873347](#).
66. Digre KB. Neuro-ophthalmology and pregnancy: what does a neuro-ophthalmologist need to know? *J Neuroophthalmol*. 2011 Dec;31(4):381-7. doi: [10.1097/WNO.0b013e31823920cb](#). PMID: [22089502](#).
67. Evcimen Y, Onur IU, Cengiz H, Yigit FU. Optical Coherence Tomography Findings in Pre-Eclampsia: A Preliminary Receiver Operating Characteristic Analysis on Choroidal Thickness for Disease Severity. *Curr Eye Res*. 2019 Aug;44(8):916-920. doi: [10.1080/02713683.2019.1600198](#). Epub 2019 Apr 15. PMID: [30983421](#).
68. Virdee J, Mollan SP. Photopsia. *Pract Neurol*. 2020 Oct;20(5):415-419. doi: [10.1136/practneurol-2019-002460](#). Epub 2020 Jun 14. PMID: [32536606](#).
69. Inan S, Polat O, Cetinkaya E, Inan UU. Bilateral serous retinal detachment accompanied by a rare intraretinal fluid configuration in preeclampsia and PRES Syndrome. *Rom J Ophthalmol*. 2019 Jan-Mar;63(1):86-90. PMID: [31198902](#); PMCID: [PMC6531768](#).
70. Panwar N, Huang P, Lee J, Keane PA, Chuan TS, Richhariya A, Teoh S, Lim TH, Agrawal R. Fundus Photography in the 21st Century—A Review of Recent Technological Advances and Their Implications for Worldwide Healthcare. *Telemed J E Health*. 2016 Mar;22(3):198-208. doi: [10.1089/tmj.2015.0068](#). Epub 2015 Aug 26. PMID: [26308281](#); PMCID: [PMC4790203](#).

71. Silverman RH, Urs R, Wapner RJ, Bearely S. Plane-Wave Ultrasound Doppler of the Eye in Preeclampsia. *Transl Vis Sci Technol*. 2020 Sep 14;9(10):14. doi: 10.1167/tvst.9.10.14. PMID: 32974086; PMCID: PMC7490228.
72. Neudorfer M, Spierer O, Goder M, Newman H, Barak S, Barak A, Asher-Landsberg I. The prevalence of retinal and optical coherence tomography findings in preeclamptic women. *Retina*. 2014 Jul;34(7):1376-83. doi: 10.1097/IAE.000000000000085. PMID: 24393833.
73. Ruia S, Tripathy K. Fluorescein Angiography. 2023 Aug 25. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: 35015403.
74. Bennett AG, Rudnicka AR, Edgar DF. Improvements on Littmann's method of determining the size of retinal features by fundus photography. *Graefes Arch Clin Exp Ophthalmol*. 1994 Jun;32(6):361-7. doi: 10.1007/BF00175988. PMID: 8082844.
75. Aissopou EK, Papathanassiou M, Nasothimiou EG, Konstantonis GD, Tentolouris N, Theodossiadi PG, Papaioannou TG, Sfrikakis PP, Protogerou AD. The Keith-Wagener-Barker and Mitchell-Wong grading systems for hypertensive retinopathy: association with target organ damage in individuals below 55 years. *J Hypertens*. 2015 Nov;33(11):2303-9. doi: 10.1097/HJH.0000000000000702. PMID: 26335430.
76. Nirupama R, Divyashree S, Janhavi P, Muthukumar SP, Ravindra PV. Preeclampsia: Pathophysiology and management. *J Gynecol Obstet Hum Reprod*. 2021 Feb;50(2):101975. doi: 10.1016/j.jogoh.2020.101975. Epub 2020 Nov 7. PMID: 33171282.
77. Chang KJ, Seow KM, Chen KH. Preeclampsia: Recent Advances in Predicting, Preventing, and Managing the Maternal and Fetal Life-Threatening Condition. *Int J Environ Res Public Health*. 2023 Feb 8;20(4):2994. doi: 10.3390/ijerph20042994. PMID: 36833689; PMCID: PMC9962022.
78. Bokuda K, Ichihara A. Preeclampsia up to date-What's going on? *Hypertens Res*. 2023 Aug;46(8):1900-1907. doi: 10.1038/s41440-023-01323-w. Epub 2023 Jun 2. PMID: 37268721; PMCID: PMC10235860.
79. Sharma DD, Chandresh NR, Javed A, Girgis P, Zeeshan M, Fatima SS, Arab TT, Gopidasan S, Daddala VC, Vaghasiya KV, Soofia A, Mylavaram M. The Management of Preeclampsia: A Comprehensive Review of Current Practices and Future Directions. *Cureus*. 2024 Jan 2;16(1):e51512. doi: 10.7759/cureus.51512. PMID: 38304688; PMCID: PMC10832549.
80. Alsamaan S, Alkhathami A, Alromaih AI, Almalki A, Alsuhbani AS, Al Ghramah AA, Nabeel M. Bilateral Exudative Retinal Detachment in Preeclampsia: A Case Report and Literature Review. *Cureus*. 2024 May 22;16(5):e60866. doi: 10.7759/cureus.60866. PMID: 38910608; PMCID: PMC11192425.
81. Shukla D, Maheshwari R, Ramchandani B, Kanungo S. Purtscher-like retinopathy with serous retinal detachment in preeclampsia of pregnancy: complications and management. *Retin Cases Brief Rep*. 2010 Fall;4(4):332-5. doi: 10.1097/ICB.0b013e3181c33389. PMID: 25390911.
82. Martinez A, Lakshminsetti M, Maharjan S, Aslam MA, Basnyat A, Kafley S, Reddy SS, Ahmed SS, Razzaq W, Adusumilli S, Khawaja UA. Beta-Blockers and Their Current Role in Maternal and Neonatal Health: A Narrative Review of the Literature. *Cureus*. 2023 Aug 24;15(8):e44043. doi: 10.7759/cureus.44043. PMID: 37746367; PMCID: PMC10517705.
83. Ibrahim A, Hussain N. Brief report: Metabolic acidosis in newborn infants following maternal use of acetazolamide during pregnancy. *J Neonatal Perinatal Med*. 2020;13(3):419-425. doi: 10.3233/NPM-190333. PMID: 31771084.
84. Razeghinejad MR. Glaucoma medications in pregnancy. *Oman J Ophthalmol*. 2018 Sep-Dec;11(3):195-199. doi: 10.4103/ojo.OJO_212_2017. PMID: 30505107; PMCID: PMC6219332.
85. Rashidian P. Roles, responsibilities, and job description of ophthalmic nurses, a universal definition is required. *Medical hypothesis, discovery & innovation in optometry*. 2022 Oct 28;3(2):55-6. 2): 55-56. doi:10.51329/mehdiptometry151.
86. Hedley PL, Hagen CM, Wilstrup C, Christiansen M. The use of artificial intelligence and machine learning methods in early pregnancy preeclampsia screening: A systematic review protocol. *PLoS One*. 2023 Apr 20;18(4):e0272465. doi: 10.1371/journal.pone.0272465. PMID: 37079505; PMCID: PMC10118197.
87. Zhou T, Gu S, Shao F, Li P, Wu Y, Xiong J, Wang B, Zhou C, Gao P, Hua X. Prediction of preeclampsia from retinal fundus images via deep learning in singleton pregnancies: a prospective cohort study. *J Hypertens*. 2024 Apr 1;42(4):701-710. doi: 10.1097/HJH.0000000000003658. Epub 2024 Jan 15. PMID: 38230614; PMCID: PMC10906188.